




THE RELATIONSHIP BETWEEN
ARTIFICIAL OXYTOCIN (PITOCIN)
USE AT BIRTH
FOR LABOR INDUCTION
OR AUGMENTATION
AND THE
PSYCHOSOCIAL FUNCTIONING
OF THREE-YEAR-OLDS


Claire L. Winstone, Ph.D.



Increasing Prevalence of
Childhood Disorders

- Attention Deficit/Hyperactivity Disorders
- Disruptive Behavior Disorders
- Attachment Disorders
- Autism Spectrum Disorders
- Sensory Processing Disorders
- Disorders of Regulation
- Learning Disabilities

Page 3




Inductions have Increased Dramatically

<u>1990</u>	<u>2005/2006</u>
9.5%	22.3%-41%

Two-thirds of inductions are now for non-medical reasons


Page 4



Pitocin Augmentation of Labor

In addition, approximately 55% of women receive Pitocin to augment labor (*Listening to Mothers, 2006*)


Page 5



Medical Indications for Induction of Labor Include:

- High blood pressure
- Premature rupture of the membranes (“premature” meaning that labor does not begin spontaneously soon after rupture)
- Maternal infection or medical problems such as diabetes mellitus, kidney disease or chronic pulmonary disease
- Suspected fetal jeopardy
- Fetal death
- Severe blood incompatibility
- Severe pre-eclampsia or toxemia
- Postdate pregnancy, where there is a proven danger to the baby (ACOG, 1999)

Page 6



Human Pitocin Research is conflicting

- 1960s, United States (Niswander)
- 1970s, Australia (McBride); USA (Friedman)
- 1980s, The Netherlands (Out & Vierhout)
- 1990s, Denmark (Sorensen)
- 1990s, Japan (Eishima)

Page 7




Measures have not been sensitive enough. . .

- Measure of adult cognitive function (Danish)
- Stanford-Binet: intelligence
- Wechsler Preschool and Primary Scale of Intelligence
- Prechtl's Neonatal Neurological Screening (2 studies)
- Brazelton Neonatal Behavioral Assessment Scale
- Porteus mazes, peg board, Graham Block Sort Test: fine and gross motor skills, conceptual ability
- "Psychomotor Development Scheme 0 – 15 months."
- Auditory and visual tests
- Physical examination
- Behavioral observation

Page 8



Neuropeptide Oxytocin Molecule




Discovery of oxytocin by physiologist
Sir Henry H. Dale, 1906

Page 9



Oxytocin	Pitocin
<ul style="list-style-type: none"> • Secreted by pituitary gland and heart • Released in pulsatile fashion, with burst firing during birth and breastfeeding • Hormone of “calm and connection” • Involved in: mother-infant attachment; parental behavior; courtship and monogamous pair bonds; and wider social relationships • Produced by baby during labor 	<ul style="list-style-type: none"> • Synthesized by pharmaceutical manufacturer • Released by IV drip in increasingly high doses until satisfactory labor pattern achieved • Does not cross blood-brain barrier, can cause increased pain and anxiety in labor • Used to start labor, to improve uterine contractions in labor and to prevent post-partum hemorrhage • Reduces newborn oxytocin levels


Page 10



Pitocin Package Insert

“Since the available data are inadequate to evaluate the benefits-to-risks considerations, Pitocin is not indicated for elective induction of labor”
(Monarch Pharmaceuticals, 2004, p. 1).


Page 11



Research into Birth and Psychosocial Functioning Has Been Based on Several Theories

- Obstetrical medications and drug addiction (imprinting theory)
- Birth trauma and suicide (imprinting theory)
- Obstetrical complications and AD/HD (hypoxia theory)
- Obstetrical complications/medications and schizophrenia and autism (oxytocin theory)

Page 12



Eishima (1992) found a significant relationship ($p < .005$) between overall obstetric conditions and overall neurological status.

Page 13



Eishima's Conclusions

“... even a small difference in low-risk obstetric conditions is related to differences in neonatal neurological status and neonatal behavior.”

Page 14



Imprinting in Animals

In the 1970s animal researchers suggest the concept of “hormonal imprinting” (Csaba, 2004, 2008)

Page 15



Imprinting in Humans

- Movement patterns (Cheek, 1986)
- Postures, movements, belief systems, relationships, artwork and language. (Emerson, 1998)
- Sequencing patterns, movements, interactions, relationships, play/behavior. (Castellino, 1997)
- Play/behavior and sequences. (Blasco, 2006)

Page 16



The Preliminary Study

- 6 clinicians working with a prenatal/perinatal frame of reference were interviewed about their perceptions of 3-year-old children born with the use of Pitocin.


Page 17



1. 6 clinicians answered 9 questions.
2. Content analysis of interviews yielded groups of items for survey.
3. 6 clinicians voted on best items to represent groups.
4. Survey developed from items.
5. Psychosocial items regrouped into "dimensions".


Page 18





The
Dissertation
Study


Page 19



A Survey on the Relationship
between Pitocin Use at Birth and
Psychosocial Functioning at Age
Three

Methodology


Page 20



The survey contained
three sets of item groups:

1. Psychosocial item groups clustered into
“dimensions”:
 - Sensory processing
 - Motor coordination
 - Affect/mood regulation
 - Self-esteem
 - ANS regulation
 - Interpersonal functioning
 - Mother-child attunement

Page 21



2. "Context of Birth" items relating to mother's experiences of pregnancy, labor, delivery and extended post-partum.

3. Demographic items.



Sample Survey Item

My child strongly resists pressure, directions, being hurried:

Never

Always



Criteria for Survey Participation

1. Mother was over 18 at the time of giving birth. Vaginal birth (not by cesarean) in an American hospital.
2. No immediately prior fetal/infant loss (miscarriage/abortion/stillbirth).
3. No street drugs or alcohol.
4. Child not premature (under 37 weeks) and/or low birth weight
5. No significant medical complications for mother or baby.
6. Mother in ongoing relationship with the father.



Why Age Three?

- More social demands on child.
- Brain is essentially fully developed—with all major fiber tracts identifiable.
- Early mother-infant attunement and resulting maturation of the right hemisphere essentially completed.
- Functioning at age three is fairly predictive of functioning at school age.

Page 25



Main Hypothesis

There is a relationship between the use of Pitocin at birth and the psychosocial functioning of children at age three.

Page 26




Online Survey Run

- The survey was run on a survey-hosting website (www.qualtrics.com) for one month.
- 514 completed surveys were analyzed.
- 16 participants were unsure if they received Pitocin or not, leaving 498 for the Pitocin-related analyses.


Page 27






Results


Page 28





Initial Data Analyses

Page 29




Non-significant Findings

Neither total score on psychosocial functioning, nor the individual dimensions (item groups) distinguished between children who did and did not receive Pitocin.


However, there was something interesting . . .

Page 30




Nine Items Significant For Pitocin

Survey Item	Dimension	Significance: 2-tailed
Reacts strongly to directions, demands	Interpersonal Dynamics: Responses to structure/support	.023
Strong need to be in control	Interpersonal Dynamics: Control/Cooperation/ Leadership	.013
Dominates other children	Interpersonal Dynamics: Control/Cooperation/ Leadership	.016
Communicates "yes" and "no" when needs to	Interpersonal Dynamics: Boundaries	.020
Mother feels she can never satisfy her child	Interpersonal Dynamics: Mother's Experience of Relationship	.048
Mother sees relationship as easy and cooperative	Interpersonal Dynamics: Mother's Experience of Relationship	.005
Rushes from one thing to next and rarely slows down	Sequencing: Beginnings	.011
Has/had a pattern of self-harming behaviors	ANS Regulation: Activity Level	.053
As an infant, had ear infections	ANS Regulation: Physiological	.038



*t test for Independent Means:
Pitocin/No Pitocin Use and Scaled Context of Birth Items*

Survey Item	t test		Sig. (2-tailed)
	t	For Equality df	
Ease of birth (Easy/Difficult)	4.971	459.955	.000***
Birth reflections (Positive/Negative)	5.353	406.685	.000***
Aspects of this birth still troubling or unresolved	2.132	469.510	.034*
Mother felt she and birth team worked well together	4.261	45.058	.000***
Others took control of birth: mother felt powerless	6.308	401.393	.000***
Felt overwhelmed, out of control, scared by labor	1.926	465.760	.055
Labor/delivery was excruciating, unbearable	3.406	441.690	.001**
Wonders if in labor baby felt disconnected, scared	4.939	408.202	.000***
Baby with Mom most of first hour post-partum	2.793	441.956	.005**




Context of Birth & Pitocin Results


Pitocin mothers: $n = 231$
no-Pitocin mothers: $n = 267$

Total score for all scaled Context of Birth items distinguished between Pitocin and no-Pitocin births to a statistically significant degree ($p = .000$).

Page 33








Factor 8: Attribute Assertiveness	Dimension in which Item was Grouped	Factor Loading
Sensitive to intrusion	Interpersonal Dynamics: Boundaries	0.2
Reacts strongly when things change quickly	Mother-Child Attunement: Transitions	0.19
Copes with transitions better when involved	Mother-Child Attunement: Transitions	0.25
Responds well to being allowed to self-pace	ANS Regulation: Timing/Pacing/Tempo	0.23
Digestive problems	ANS Regulation: Physiological	-0.16
Picky eater	ANS Regulation: Physiological	-0.55
Problems with sleeping	ANS Regulation: Physiological	-0.18
Baby ear infections	ANS Regulation: Physiological	-0.42
Baby colicky, fussy, hard to soothe	ANS Regulation: Physiological	-0.25

Page 36

Means for factor 8: Assertiveness

Pitocin group (n = 231) 1.346381
 No Pitocin group (n = 267) 1.858906



Page 37



Factor 12: Attribute: Need to Control Environment	Dimension	Factor Loading
Craves firm pressure	Sensory Processing: Tactile/Proprioceptive	0.32
Procrastinates on starting things	Sequencing	0.3
Loses focus in the middle	Sequencing	0.15
Responds well to self-pace	ANS Regulation: Timing/Pacing/Tempo	0.16
Baby colicky, fussy, hard to soothe	ANS Regulation: Physiological	0.23
Copes with transitions better when involved	Mother-Child Attunement: Transitions	0.26
Resists receiving help from others	Interpersonal Dynamics: Responses to structure/support	0.15
Strongly resists pressure; being hurried	Interpersonal Dynamics: Responses to structure/support	0.18
Strong need to be in control	Interpersonal Dynamics: Control/Cooperation/Leadership	0.33
Aggressive	ANS Regulation: Activity Level	-0.19
Digestive problems	ANS Regulation: Physiological	-0.25
Baby hypersensitive to movement/position	Sensory Processing: Vestibular	-0.23
Reacts strongly when things change quickly	Mother-Child Attunement: Transitions	-0.18
Difficulty with others' timing/transitions	Interpersonal Dynamics: Responses to structure/support	-0.18
Gets upset when things don't go their way	Interpersonal Dynamics: Control/Cooperation/Leadership	-0.16



Means for Factor 12: Need to Control Environment



Pitocin group (n = 231) 2.481183
 No Pitocin group (n = 267) 2.169549



Page 39



Relationship of Assertiveness to Context Score and Pitocin Use

Context scores  = Child's Assertiveness 

The more negative/stressful the Context of Birth for mother, the more the child experiences internal dysregulation and needs to use Assertiveness to get help with dysregulation

Pitocin use  = Child's ability to employ assertiveness 

Page 40



Significant Difference between Exclusive Breastfeeding and Other Forms of Feeding with Pitocin and No Pitocin at Birth

	Baby Fed Entirely at Breast	Combination Breast/Bottle-feeding Plus Exclusive Bottle-feeding	Totals
Pitocin	96	135	231
No Pitocin	231	16	247
Total	327	151	478

Fisher's Exact Test: (95% CI, $p < .000$)

Page 41





Discussion



Eishima's Study of Newborns

“... even a small difference in low-risk obstetric conditions is related to differences in neonatal neurological status and neonatal behavior.”

Page 43



This study supports Eishima's Findings Through Age Three

Factor 8: Assertiveness

Factor 12: Need to Control Environment

Both suggest themes of difficulties in:

- Interpersonal dynamics
- Mother-child attunement
- Autonomic nervous system regulation
- Sensory processing
- Sequencing

Page 44



Five Infant Items

- Of the five items in the survey describing characteristics of the child in infancy, groups of two or more appeared in majority of the PCA factors studied (7 of the 12 studied), including the two found significant for Pitocin:
- *Suggests a continuity of disturbance in functioning from infancy to age three.*
- Faulty oxytocin imprinting could explain this continuity.

Page 45



The Polyvagal Theory of Stephen Porges

Page 46




Jackson's Theory of Dissolution

"The higher nervous system arrangements inhibit (or control) the lower, and thus, when the higher are suddenly rendered functionless, the lower rise in activity."



—John Hughlings Jackson (1835-1911)
Father of English Neurology
Quoted by Stephen Porges 11/01



We play our **newest, best card** first, if that doesn't work (or has not worked in the past as determined by the amygdala), we try our **older, second card**. If that doesn't work, we play our **oldest, last card**. If that doesn't work we are in extreme danger of death.

Slide prepared by John Chitty, Colorado School of Energy Studies, www.energyschool.com

“Assertiveness”
(decreased by Pitocin) ≈
Social Engagement

“Need to Control Environment”
(increased by Pitocin) ≈
Sympathetic Activation (Fight/flight)

Page 48



Oxytocin Receptors

- During pregnancy in humans, oxytocin receptor numbers increase in the uterus: beginning at six times the number in a non-pregnant uterus, up to 80-fold at term (Fuchs & Fuchs, 1984)

Page 49



Hormonal Imprinting

“Excess of the target hormones . . . which are able to bind to the receptors, provoke faulty imprinting in the critical periods with life-long morphological, biochemical, functional or behavioural consequences.”
(Csaba, 2008, p. 1)

Page 50



- PCA findings suggest that the binding of exogenous oxytocin (Pitocin) to oxytocin receptors during labor and delivery may result in faulty imprinting affecting aspects of ANS regulation and interpersonal dynamics.
- If both mother and baby experience faulty oxytocin imprinting, both may be contributing to a relative lack of attunement in the relationship, as reflected in significant differences in scores.

Page 51



Major Findings of the Study

Significant relationships found between Pitocin and:

- Aspects of psychosocial functioning at age three, particularly ANS functioning and the resulting interpersonal dynamics.
- Reduced time spent together by mother and baby in first hour post-partum.
- Reduced exclusive feeding at the breast in the first six months.

Page 52



Conclusions

- Pitocin use at birth appears to have psychosocial, and probably neurobiological, consequences for mother and child.
- Little is known about the long-term consequences of interfering with the hormonal physiology of childbirth in general and with the oxytocin system in particular.
- Increased Pitocin use at birth may be a contributing factor to the concurrent increase in a variety of disorders of regulation in children.

Page 53



Conclusions, cont'd

- Further research needed to expand on present study and extend the investigation to the psychosocial functioning of older children.
- Possibility that other obstetrical interventions may also have long-term effects that require study.
- Encourage re-examination of Pitocin use in birth, and careful weighing of the risks and benefits of its use, for the benefit of the most vulnerable.

Page 54

